

Name: _____ Total Points: _____
(Last) (First)

Physics 201

Exam 3

Write also your name in the
appropriate box of the scantron


Multiple choice questions

Answer all of the following questions. Read each question carefully. **Fill the correct bubble on your scantron sheet.** Each question has exactly one correct answer. All questions are worth the same amount of points.

1. A point mass m has kinetic energy KE_1 measured in an inertial frame R_1 and kinetic energy KE_2 measured in another inertial frame R_2 . The velocity of R_2 with respect to R_1 is $\vec{v} \neq 0$. Which of the following is true for sure?

- A. $KE_2 = KE_1$
- B. $KE_2 = KE_1 + 1/2mv^2$
- C. $KE_2 < KE_1$
- D. $KE_2 > KE_1$
- E. $KE_2 \neq KE_1$


2. The center of mass of the system of particles shown in the diagram is at point



- A. a
- B. b
- C. c
- D. d
- E. e

Name: _____ Total Points: _____
(Last) (First)

3. Using a mallet, you strike a ball of mass 0.50 kg that is initially at rest. The force F on the ball as a function of time is plotted in the figure. At $t=2.0\text{ms}$, the speed of the ball is



- A. 10 m/s
- B. 8.0 m/s
- C. 6.0 m/s
- D. 4.0 m/s
- E. 2.0 m/s

Questions 4 through 10 all refer to the same problem.


A block of mass $m=0.5\text{kg}$ is initially at rest on a frictionless 30° incline as shown on the figure below. The block is resting against a massless spring of spring constant $k=100\text{N/m}$. The spring is not attached to the block.

Displacements are measured along an x -axis directed up along the incline.

Take the origin $x=0$ to be the location where the spring is not compressed.

Since the spring is not attached to the block, the spring loses contact with the block for $x>0$.

Take $g=10\text{ m/s}^2$.



4. What is the location x_{eq} of the block at equilibrium?

- A. -2.5 cm
- B. -4.33 cm
- C. -5 cm
- D. -7 cm
- E. -10 cm

The block is displaced by an additional 4 cm down the incline from its equilibrium position and then released (in other words, the block is released at $x=x_{\text{eq}}-4\text{cm}$ with no initial velocity).

In what follows, $U(x)$ is the potential energy of the system block+spring+Earth at position x along the incline.


5. How do $U(x_{\text{eq}}-4\text{cm})$ and $U(x_{\text{eq}})$ compare?
- A. $U(x_{\text{eq}}-4\text{cm}) > U(x_{\text{eq}})$
 - B. $U(x_{\text{eq}}-4\text{cm}) < U(x_{\text{eq}})$
 - C. $U(x_{\text{eq}}-4\text{cm}) = U(x_{\text{eq}})$
 - D. Can't tell. There is not enough information.
6. After its release, as the block travels up the incline, what can you say about the total mechanical energy E_{mec} of the system block+spring+Earth?
- A. E_{mec} is constant.
 - B. E_{mec} decreases as x increases.
 - C. E_{mec} increases as x increases.
 - D. From $x=x_{\text{eq}}-4$ cm to $x=0$ cm, E_{mec} increases. Then E_{mec} decreases as x increases.
 - E. From $x=x_{\text{eq}}-4$ cm to $x=0$ cm, E_{mec} decreases. Then E_{mec} increases as x increases.
7. After its release, what is the position x_{max} of the block up the incline when it comes momentarily to rest for the first time?
- A. -1.5 cm
 - B. 1.5 cm
 - C. 1.95 cm
 - D. 4 cm
 - E. 10 cm
8. How do $U(x_{\text{eq}}-4\text{cm})$ and $U(x_{\text{max}})$ compare?
- A. $U(x_{\text{eq}}-4\text{cm}) > U(x_{\text{max}})$
 - B. $U(x_{\text{eq}}-4\text{cm}) < U(x_{\text{max}})$
 - C. $U(x_{\text{eq}}-4\text{cm}) = U(x_{\text{max}})$
 - D. Can't tell. There is not enough information.

Name: _____ Total Points: _____
(Last) (First)

9. As the block travels up the incline, where is its kinetic energy maximum?
- A. at x_{\max}
 - B. at x_{eq}
 - C. at the origin $x=0$
 - D. at some location between $x=0$ and x_{\max}
 - E. at some location between $x_{\text{eq}}-4\text{cm}$ and x_{eq}
10. As the block travels up the incline, where is the magnitude of its acceleration maximum?
- A. at x_{\max}
 - B. at x_{eq}
 - C. at the origin $x=0$
 - D. at $x_{\text{eq}}-4\text{cm}$
 - E. at some location between $x_{\text{eq}}-4\text{cm}$ and x_{eq}

Questions 11 through 17 all refer to the same problem.

On a frictionless horizontal track, block A of mass 3 kg collides with block B of mass 2 kg. Before the collision, the velocity of A is $\vec{v}_A = 3\hat{x} \text{ m/s}$ and the velocity of B is $\vec{v}_B = 2\hat{x} \text{ m/s}$ as indicated on the figure below. After the collision, it is observed that A and B have the same velocity \vec{v} .



11. During the collision, what is the direction of the net force on A?
- A. \hat{x}
 - B. $-\hat{x}$
 - C. Undefined, since the net force is 0
12. During the collision, what can you say about the magnitude $|\vec{p}_A|$ of the momentum of A?
- A. $|\vec{p}_A|$ increases
 - B. $|\vec{p}_A|$ decreases
 - C. $|\vec{p}_A|$ stays the same
 - D. Can't say anything. There is not enough information.

Name: _____ Total Points: _____
(Last) (First)

13. What is the velocity of the center of mass of the system A+B before the collision (in m/s)?

- A. 0
- B. $2\hat{x}$
- C. $2.5\hat{x}$
- D. $2.6\hat{x}$
- E. $5\hat{x}$

14. What is the velocity \vec{v} of A and B after the collision (in m/s)?

- A. 0
- B. $2\hat{x}$
- C. $2.5\hat{x}$
- D. $2.6\hat{x}$
- E. $5\hat{x}$

15. In the center of mass frame, what is the kinetic energy of the system before the collision (in Joules)?

- A. 0
- B. 0.6
- C. 1.2
- D. 17.5
- E. 35

16. In the center of mass frame, what is the kinetic energy of the system after the collision (in Joules)?

- A. 0
- B. 0.6
- C. 1.2
- D. 17.5
- E. 35

Name: _____ Total Points: _____
(Last) (First)

17. Is the collision an elastic collision?

- A. No in all inertial frames.
- B. Yes in all inertial frames.
- C. Yes in the center of mass frame. No in an inertial frame fixed with respect to the horizontal track.
- D. No in the center of mass frame. Yes in an inertial frame fixed with respect to the horizontal track.
- E. Can't tell. There is not enough information. N